

Production of Xylanase and β -Xylosidase by *Aspergillus niger* NRC 107

Abdel-Naby, Mohamed A. and Dae Young Kwon^{1*}

Department of Chemistry of Natural and Microbial Products,
National Research Center, Dokki, Cairo, Egypt

¹Food Chemistry Laboratory, Korea Food Research Institute,
San 46-1, Paekhyondong, Poondang, Songnam 463-420, Korea

Aspergillus niger NRC 107에서의 Xylanase와 β -Xylosidase의 생산

압델나비 모하메드 · 권대영^{1*}

도끼국립연구소 천연·미생물자원 화학과, ¹한국식품개발연구원 화학연구소

Abstract— The production of xylanase and β -xylosidase was investigated in submerged culture of *Aspergillus niger* NRC 107. The maximum production occurred when the pH was controlled at 6.0 during the fermentation. Among the various carbon sources investigated, corn-cob xylan (1.5%, w/v) yielded maximal production of the enzymes. The NaNO₃ was the most favorable nitrogen source for enzyme production and KH₂PO₄ concentration at 0.3%(w/v) was found to be optimum. Incorporation of wheat bran to the culture medium improved xylanase production. Addition of L(-) sorbose to the culture medium promoted the secretion of β -xylosidase. It was possible to increase the production of xylanase (39.43 units/ml) and that of β -xylosidase (4.2 units/ml) by submerged culturing the *A. niger* NRC 107 in the modified medium.

Hemicellulose is one of the major components of lignocellulosic materials comprising 15 to 35% of agricultural and forestry residues. Xylans (β -1,4-D-xylose polymer) are the major components of hemicellulose (1). Extensive degradation of xylans to monosaccharides is achieved by cooperative action of multi-enzymes; endo- β -xylanase, exo- β -xylanase and β -xylosidase (2). In general, endo- β -xylanase (usually called xylanase, E.C. 3.2.1.8) randomly cleaves the 1,4- β -xylosidic linkages in xylans and exo- β -xylanase and β -glucosidase hydrolyze the 1,4- β -D-xylans or xylooligosaccharides so as to remove successive D-xylose residue from the non-reducing termini. β -Xylosidase (E.C. 3.2.1.37) prefer to short xylooligosaccharides. Many microbial sources have been reported to produce these enzymes (3-8).

The cost of production and low yield of these

Key words: Hemicellulose, xylanase, β -xylosidase, xylanase fermentation

*Corresponding author

enzymes are the major problems for practical application (9). Therefore, the availability of highly active xylanase would be of great significance. This requires the selection of suitable strain and the development of fermentation technology for producing the enzymes in quantity.

This study was undertaken to investigate some factors affecting the production of xylanases and β -xylosidase by a local isolate *Aspergillus niger* NRC 107. From a certain comparison made with other fungi, it is apparent that this culture produces large amount of xylanases. In addition, *A. niger* NRC 107 secretes high amounts of β -xylosidase.

Materials and Methods

Microbes and Materials

The microorganisms utilized in the present study were obtained from the fungal collection of the Centre of Cultures of National Research Centre, Cairo,

Egypt. They were maintained on potato dextrose agar slants at 25°C and subcultured twice a month.

Oat spelt xylan was obtained from Fluka, Switzerland and *p*-nitrophenol- β -D-xylosidase was purchased from Sigma Chemical Co., MO (U.S.A). Corn-cob xylan and wheat straw xylan were prepared by the method of Whistler *et al.* (10).

Media and culture conditions

The basal salt medium for the growth of the fungal and the enzyme production was that of Mandels and Stenrnberg (11). The medium contained in each liter: KH_2PO_4 , 2.0g; $(\text{NH}_4)_2\text{SO}_4$, 1.4g; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.3g; CaCl_2 , 0.3g; urea, 0.3g; Tween-80, 1 ml; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 5.0 mg; $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 1.6 mg; $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 1.4 mg; CoCl_2 , 2.0 mg; and corn-cob xylan, 10g. The pH was adjusted to 5.0. The addition of different substrates to the medium is indicated in each experiment. Cultivation was made in 250 ml Erlenmeyer flasks each containing 50 ml of sterilized media. One milliliter of spore suspension (8×10^7 spores/ml) obtained from 7-day mother culture was used for inoculation. Cultivation was performed at 30°C on a rotary shaker (180 r.p.m.). The cultures were harvested on the 7th day of growth by filtration through glass wool filter and then centrifuged. The clear supernatants were used for enzyme assays.

The pH control was done by adding either 1 N NaOH or 1 N HCl every 12 hr.

Determination of enzyme activities

Xylanase activity in the culture filtrate was determined from the amount of the reducing sugars formed in terms of xylose according to the method of Somogyi (12). The half milliliter of appropriately diluted culture filtrate was added to 0.5 ml of 1.0% (w/v) xylan in 0.05 M phosphate buffer (pH 5.0). The reaction mixture was incubated at 45°C for 30 min. One unit (U) of xylanase activity was defined as the amount of enzyme liberating one mole of reducing sugars as a xylose per min.

β -Xylosidase activity was also assayed by measuring the amount of *p*-nitrophenol liberated from *p*-nitrophenyl- β -D-xyloside (PNPX) (13). The assay mixture consisted of 0.5 ml of 0.1 M phosphate buf-

fer, pH 5.0, 0.5 ml of 0.05 M PNPX and 1 ml of enzyme solution. The reaction mixture was incubated at 40°C for 15 min, then 2 ml of 1M Na_2CO_3 were added to stop the reaction. The color developed was read at 400 nm with the spectrophotometer. One unit (U) of β -xylosidase activity was defined as the amount of enzyme that produced one mole of *p*-nitrophenol per min.

Protein content of mycelium

The protein content in the mycelium was determined indirectly by estimation of Kjeldahl nitrogen of total solids (14). With determined nitrogen content (N), mycelium protein was calculated as $N \times 6.25$. Soluble protein in culture filtrates was determined by the method of Lowry *et al.* (15).

The results shown represent the means of at least three separately undertaken experiments.

Results and Discussion

Screening of some fungal strains for the production of extracellular enzymes

All tested fungi were grown on the basal medium containing corn-cob xylan. The culture filtrates were investigated for extracellular xylanase and β -xylosidase activities on the 7th day of growth (Table 1). *Aspergillus niger* NRC 107 was found to be the most potent fungus for xylanase and β -xylosidase production, followed by *A. oryzae* NRC 13. On the other hand, *Fusarium oxysporum* 3A was comparatively the lowest xylanase producer. Therefore, *A. niger* NRC 107 was selected for further works.

Optimization of *A. niger* NRC 107 culture conditions for xylanase and β -xylosidase

Effect of pH of culture medium: The pH of culture medium was maintained at pH 3.0, 4.0, 5.0, 6.0 and 7.0 through the fermentation process (Fig. 1). Maximal enzyme production was detected at pH 6.0. These results coincide with those reported for xylanase and β -xylosidase production by *A. terreus* IJIRA (16) and *A. terreus* 603 (6). At pH 6.0, xylanase activity was 4.5-fold that at pH 3.0. The most significant effect of low pH was demonstrated by the complete inhibition of β -xylosidase at pH 3.0. The low enz-

Table 1. Survey of fungal cultures for the production of xylanase and β -xylosidase

Microorganisms	Final pH of C.F.*	Protein content of		Enzyme activities (unit/ml)	
		mycelium (mg/50 ml culture)	C.F. (mg/ml)	Xylanase	β -Xylosidase
<i>Aspergillus niger</i> NRC107	7.5	70.3	0.65	6.84	0.54
<i>Aspergillus oryzae</i> NRC13	8.5	84.3	0.54	4.64	0.19
<i>Aspergillus saitoi</i> B22	7.0	60.3	0.49	1.29	0.03
<i>Fusarium oxysporum</i> 3A	7.4	43.2	0.66	0.39	0.00
<i>Penicillium citrinum</i> PY30	6.8	90.3	0.55	2.1	0.12
<i>Penicillium funiculosum</i> WD7	7.6	36.3	0.44	0.88	0.04
<i>Trichoderma viride</i> 250	6.5	55.3	0.74	2.57	0.07
<i>Trichoderma harzianum</i> 27	6.8	64.3	0.39	1.84	0.00

*C.F. is the abbreviation of culture filtrate. This is the same as in this and following Tables and Figures.




Fig. 1. Effect of pH on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Protein content of mycelium (■), protein content of culture filtrate (□), xylanase (●), and β -xylosidase (○). Standard medium: corn-cob xylan, 10 g/l; $(\text{NH}_4)_2\text{SO}_4$, 1.4 g/l.

yme activity at pH 3.0 may at least partially be due to inactivation of the enzyme at this pH rather than to inhibition of enzyme biosynthesis (17). This view may explain the production of appreciable amounts of extracellular protein at pH 3.0 and the large increase of enzyme yield with limited increase of extracellular protein and mycelial biomass at higher pH's. Therefore pH 6.0 was used.

Table 2. Effect of different carbon sources on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Carbon source*	Enzyme activities(unit/ml)	
	Xylanase	β -Xylosidase
Control	8.24	0.68
Cellulose	1.14	0.00
Starch	5.94	0.22
Maltose	3.14	0.06
Sucrose	2.44	0.03
Lactose	6.24	0.39
Glucose	3.69	0.00
Xylose	3.29	0.00
Arabinose	1.89	0.04

*Carbon levels of each samples were kept constant equivalent to the amount of the control: 10 g/l corn-cob xylan.

Effect of carbon sources: The production of xylanase and β -xylosidase was compared when *A. niger* NRC 107 was grown on the basal media containing different carbon sources (Table 2). The carbon level was kept constant (equivalent to 1.0% corn-cob xylan). The results showed that xylan served as a superior carbon source for xylanase and β -xylosidase production by *A. niger* NRC 107. This results were in agreement with those reported for xylanase and β -xylosidase from *Trichoderma viride* (18), *A. terreus* (16), *Talaromyces byssochlamydoides* (19) and *T. lignorum* (20). On the other hand, feeble production of xylanase and β -xylosidase were obtained

with glucose, maltose, xylose and lactose. The utilization of these sugars in the culture medium indicated good cell growth (data not shown). These results were in accordance with those of reported data (3, 7, 20, 21). Many investigators found a considerable yield of xylanase and β -xylosidase on using a culture medium containing cellulose as a unique carbon source (16, 20, 22). In *A. niger* NRC 107, however, a feeble yield (15% of that with xylan) of enzyme was observed upon using cellulose as a sole carbon source. This may be partially due to poor biomass production, perhaps because of the lack of cellulase to break down the cellulose to soluble sugars. In fact, protein content of mycelium with cellulose (22.13 mg/50 ml culture) was about 30% of that with xylan (68.94 mg/50ml culture).

In further studies on the effect of carbon source on enzyme production, different xylan substrates at various concentrations were investigated for production of xylanase and β -xylosidase from *A. niger* NRC 107 (Table 3). Of the substrates investigated corn-cob xylan was the most favorable for enzyme production. Maximal enzyme yield was observed at 15 g/l corn-cob xylan.

Effect of nitrogen sources: On equivalent nitrogen basis, the nitrogen sources from the basal medium ((NH₄)₂SO₄ + urea) were substituted by several

kinds of nitrogen sources (Table 4). Of the nitrogen sources investigated NaNO₃ was the most favorable for the production of active xylanase and β -xylosidase from *A. niger* NRC 107. This result was similar to that reported by Ismail *et al.* (6). Ammonium compounds such as NH₄NO₃, (NH₄)₂SO₄

Table 4. Effect of different nitrogen sources on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Nitrogen source*	Enzyme activities(unit/ml)	
	Xylanase	β -Xylosidase
Control	12.2	1.07
NaNO ₃	17.8	1.85
(NH ₄)Cl	11.3	1.88
(NH ₄)NO ₃	10.9	1.6
(NH ₄) ₂ SO ₄	13.9	1.03
Urea	7.2	0.44
Meat extract	2.8	0.06
Peptone	4.7	0.74
Yeast extract	3.9	0.54
Casein hydrolyzate	2.9	0.37

*Nitrogen levels of each samples were kept constant equivalent to the amount of the control: (NH₄)₂SO₄, 1.4 g/l + urea, 0.3 g/l. Carbon source (corn-cob xylan) levels of each samples were fixed to concentration: 15 g/l, which was optimized from the Table 3.

Table 3. Effect of various xylan substrates on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Xylan substrates	Xylan concentration (g/l)	Protein content of		Enzyme activities (unit/ml)	
		mycelium (mg/50 ml culture)	C.F. (mg/ml)	Xylanase	β -Xylosidase
Corn-cob	5	60.3	0.68	4.39	0.45
	10	68.9	0.77	8.24	0.68
	15	82.2	1.05	12.24	1.07
	20	94.7	1.24	10.24	0.84
Wheat straw	5	53.2	0.74	3.69	0.39
	10	60.8	0.79	7.74	0.7
	15	71.2	0.84	9.28	0.84
	20	78.5	0.93	8.88	0.9
Oat splot	5	66.4	0.82	6.69	0.44
	10	73.2	0.91	10.33	0.63
	15	88.2	0.97	9.24	0.71
	20	94.3	1.11	8.21	0.71

Nitrogen levels of each samples were kept constant equivalent to the amount of the control: (NH₄)₂SO₄, 1.4 g/l + urea, 0.3 g/l.

and NH_4Cl showed moderate levels of xylanase. On the other hand, organic nitrogen sources (meat extract, peptone, yeast extract, casein hydrolyzate and urea) were relatively unsuitable for xylanase and β -xylosidase production from *A. niger* NRC 107. These results were in contrast to that reported by Ghash and Kundu (16).

Effect of phosphate level: The effect of phosphate level on the production of xylanase and β -xylosidase was investigated using different concentrations of KH_2PO_4 (Fig. 2a). The results showed that KH_2PO_4 concentration at 0.3% (w/v) was the most favorable concentration for the production of both extracellular enzymes. Higher levels had an adverse effect on xylanase production.

Effect of surfactants: The addition of surfactants was found to increase extracellular enzyme activity for a variety of microorganisms (23). The effect of surfactants on production of xylanase and β -xylosidase from *A. niger* NRC 107 was investigated using different concentration of Tween-80 (Fig. 2b). As shown in the results, the increase of Tween-80 concentration from 0.1% to 0.3% led to activity increase in xylanase (by 16.5%) and β -xylosidase (by 13.0%) relative to control. Higher concentrations, however, had an adverse effect on the enzyme production. These results were similar to those of Szczodrak *et al.* (24). On the other hand, Ghash and Kundu (16) found no significant effect of Tween-80 and Triton X-100 on xylanase and β -xylosidase production by *A. terreus*.

Effect of addition of organic substrates: The effect of various organic substrates and inducers on the production of xylanase and β -xylosidase from *A. niger* NRC 107 was investigated as shown in Table 5. Corn steep had an adverse effect on xylanase production. On the other hand, remarkable increase (74%) in β -xylosidase was shown by the addition of L(-) sorbose at 0.3% (w/v). Bisaria *et al.* (25) found that L(-) sorbose at 0.5% (w/v) improved β -glucosidase yield from *Trichoderma reesei* QM 9414. Table 5 shows that rice bran at 1.0% (w/v) provided a slight increase in xylanase activity while wheat bran was the best for xylanase. Thus addition of 1.5% (w/v) wheat bran to the culture medium




Fig. 2. Effect of phosphate (a) and Tween-80 (b) concentrations on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Standard medium: Corn-cob xylan, 15 g/l; NaNO_3 , 2.67 g/l was used. The concentration of Tween-80 in the phosphate effect testing (Fig. 2a) and the concentration of KH_2PO_4 in Fig. 2b were 1 ml/l, and 3.0 g/l, respectively. Symbols: xylanase (●), and β -xylosidase (○).

resulted in 55.3% and 36.0% increase in xylanase and β -xylosidase respectively. This result coincides with those reported by other investigators (18, 19, 26 and 27). The highest yield of xylanase (39.43 unit/ml) and a high one β -xylosidase (4.2 units/ml) were achieved upon addition of both wheat bran (at 1.5%, w/v) which is the best for xylanase production, and L(-) sorbose (at 0.3%, w/v) which is the best for β -xylosidase (Table 5).

Table 5. Effect of some organic substrates on the production of xylanase and β -xylosidase by *A. niger* NRC 107.

Organic substrates	Substrate concentration (g/l)	Protein content of		Enzyme activities (unit/ml)	
		mycelium (mg/50 ml culture)	C.F. (mg/ml)	Xylanase	β -Xylosidase
None*	0	98.7	1.39	26.7	2.94
Corn-steep	0.5	156.7	1.89	20.4	2.2
	1.0	192.5	2.56	18.7	2.08
	1.5	214.3	2.56	18.7	1.89
Rice bran	5	110.3	1.58	25.8	2.7
	10	139.7	1.69	27.7	2.78
	15	156.8	1.84	26.6	2.16
Wheat bran	10	102.3	1.47	34.6	3.14
	15	118.9	1.66	38.7	3.54
	20	134.3	1.73	37.7	3.5
L(-) sorbose	1	109.3	1.42	28.9	3.41
	3	121.3	1.54	29.6	4.52
	5	133.7	1.63	26.3	4.31
Wheat bran+	15				
L(-) sorbose	3	144.7	1.72	39.4	4.2

*For the experimentals, major medium composition of control is as follows: corn-cob xylan, 15 g/l; NaNO_3 , 2.67 g/l; KH_2PO_4 , 3.0 g/l, Tween-80, 0.3 (%v/v)

Production of xylanase and β -xylosidase in optimized medium

According to the above mentioned results a medium was formulated taking into consideration the most favorable conditions reached. The each liter of medium consisted of: corn-cob xylan, 15g; NaNO_3 , 2.67g; KH_2PO_4 , 3.0g; CaCl_2 , 0.3g; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.3 g; wheat bran, 15g; L(-) sorbose, 3.0g; Tween 80, 3 ml; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 5.0 mg; $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 1.6 mg; $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$, 1.4 mg; and CoCl_2 , 2.0 mg. The pH was controlled through the fermentation processes of 7 days at pH 6.0.

With this optimized media, the course of production of xylanase, β -xylosidase, extracellular protein and mycelial biomass were investigated as shown in Fig. 4. After inoculation there was a lag of 36 hr before extracellular protein and enzymes reached detectable levels. Extracellular xylanase appeared earlier on the second day of growth. The appearance of β -xylosidase activity was lagged approximately by 24 hr behind xylanase production. Xylanase and β -xylosidase were apparently produced during active growth and there is a close linear relationship between enzyme formation and the in-




Fig. 3. Fermentation profile for xylanase and β -xylosidase production by *A. niger* NRC 107 in optimized medium.

Corn-cob xylan, 15 g/l; NaNO_3 , 2.67 g/l; KH_2PO_4 , 3.0 g/l; Tween 80, 3 ml/l; wheat bran, 15.0 g/l; L(-) sorbose, 3.0 g/l. Symbols: protein content of mycelium (■), protein content of culture filtrate (□), xylanase (●), and β -xylosidase (○).

crease of extracellular protein. These results were similar to those reported on the course of xylanase and β -xylosidase from other microbial sources (7, 13, 19, 24 and 28). In addition, the culture filtrate of *A. niger* 107 showed no cellulase activity (data not shown).

A. niger NRC 107 in its optimum medium afforded 39.43 unit/ml of xylanase and 4.2 unit/ml of β -xylosidase. The optimum culture medium provided 5.76-fold and 7.7-fold increases in xylanase and β -xylosidase, respectively, relative to the original basal medium (Table 1). *A. niger* NRC 107 xylanase activity was higher than those reported for *A. terreus* 603 (6), *A. wentii* Pt 2804 (29), *Talaromyces byssochloamydoides* YH-50 (19), and *Trichoderma reesei* D 1-6 (28). At the same time, the levels of β -xylosidase obtained from *A. niger* NRC 107 are higher than those obtained from *A. terreus* (16), *Trichoderma reesei* QM 9414 (13), *Penicillium funiculosum* (30) and *Streptomyces* sp. (7).

요 약

Hemicellulose를 효소에 의하여 분해시키기 위하여 *Aspergillus niger* NRC 107로부터 xylanase와 β -xylosidase의 생산 조건에 대하여 조사하였다. 이들 효소 생산 최적 pH는 6.0이었으며 탄소원 중 corn-cob xylan과 질소원 중 Na_2NO_3 이 효소 생산에 제일 좋은 기질이었으며 이들의 최적농도는 각각 15 g/l와 2.67 g/l이다. 이들 효소는 인산염(KH_2PO_4)과 Tween-80의 농도를 조절하므로 수율을 높일 수 있으며 wheat bran은 xylanase의 생산에 L(-) sorbose는 β -xylosidase의 생산에 좋은 영향을 주었다. 이와같이 이 *Aspergillus niger* NRC 107의 회분식 배양에서 이들 효소에 영향을 미치는 생산조건을 최적화한 결과 xylanase는 39.43 units/ml, β -xylosidase는 4.2 units/ml까지 생산할 수 있었다.

References

- Whistler, R.L. and E.L. Richards. 1970. Xylans, pp. 447-469. In W. Pigman, and D. Horton(ed.), *Hemicellulose in the Carbohydrates-Chemistry and Biochemistry*, Academic Press, New York.
- Frederick, M.M., C.-H. Kiang, J.R. Frederick and P.J. Reilly. 1985. Purification and characterization of endoxylanases from *Aspergillus niger*. I. Two isozymes active on xylan backbones near branched points. *Biotechnol. Bioeng.* **27**: 525-532.
- Esteban, R., J.R. Villanueva and T.G. Villa. 1982. β -Xylanases of *Bacillus circulans* WI-12. *Can J. Microbiol.* **28**: 733-739.
- Mishra, C., S. Keskar and M. Rao. 1984. Production and properties of extracellular endoxylanase from *Neurospora crassa*. *Appl. Environ. Microbiol.* **48**: 224-228.
- Mukherjee, M. and S. Sengupta. 1985. An inducible xylanase of the mushroom *Termitomyces clypeatus* differing from the xylanase/amylase produced in dextrin medium. *J. Gen. Microbiol.* **131**: 1881-1985.
- Ismail, A.S., M.A. Abdel-Naby and A.F. Abdel-Fattah. 1986. Production of hemicellulytic enzymes by fungi. *Agric. waste* **18**: 283-291.
- Nakanishi, K., K. Yokotsuka and T. Yasui. 1987. Induction of membrane-bound xylosidase in a *Streptomyces* sp. *J. Ferment. Technol.* **65**: 1-6.
- Mountfort, D.O. and R.A. Asher. 1989. Production of xylanase by the ruminal anaerobic fungus *Neocallimastix frontalis*. *Appl. Environ. Microbiol.* **55**: 1016-1022.
- Ailts, B., D. Englund and L. Brooks. 1981. Conversion of agricultural biomass to glucose for ethanol production determination of minimum equipment required and production cost, pp. 359-367. In P.F. Bente(ed.), *International Bioenergy Directory*. The Bioenergy Council, Washington, D.C..
- Whistler, R.L., J. Bachrach D.R. Bowman. 1948. Preparation and properties of corn cobs hollocellulose. *Arch. Biochem.* **19**: 25-33.
- Mandels, M. and D. Sternberg. 1976. Recent advances in cellulase technology. *J. Ferment. Technol.* **54**: 267-286.
- Somogyi, A.M. 1952. A new reagent for determination of sugar. *J. Biol. Chem.* **195**: 19-24.
- Dekker, R.F.H. 1983. Bioconversion of hemicellulose: Aspects of hemicellulase production by *Trichoderma reesei* QM 9414 and enzymatic saccharification of hemicellulose. *Biotechnol. Bioeng.* **25**: 1127-1146.
- Herbert, D., P.J. Phipps and R.E. Strange. 1971. Chemical analysis of microbial cells, pp. 209-343. In Norris, J. R. and D.W. Ribbons(ed.), *Methods in Microbiology*, pp. 1361-9. 5B, Academic Press, London.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.L. Randall. 1951. Protein measurement with Folin-phenol reagent. *J. Biol. Chem.* **193**: 265-276.

16. Ghosh, B.S. and A.B. Kundu. 1980. Induction of cellulases and hemicellulases by Tamarind "*Tamarinds indica*" kernel polysaccharide. *J. Ferment. Technol.* **58**: 135-141.
17. Bae, S.-H. and Y.-J. Choi. 1991. Purification and characterization of an extracellular xylanase of *Bacillus stearothermophilus*. *Kor. J. Appl. Microbiol. Biotechnol.* **19**: 592-597.
18. Hashimoto, S., T. Muramatsu and M. Funatsu. 1971. Studies on xylanase from *Trichoderma viride* part 1. Isolation and some properties of crystalline xylanase. *Agric. Biol. Chem.* **35**: 501-508.
19. Yoshioka, H., S. Chavanich, S. N. Nilubol and S. Hayashida. 1981. Production and characterization of thermostable xylanase from *Talaromyces byssochlamydoides* YH-50. *Agric. Biol. Chem.* **45**: 579-586.
20. Defaye, J., H. Driguez, M. John, J. Schmidt and E. Ohleyer. 1985. Induction of D-xylan degrading enzymes in *Trichoderma lignorum* by nonmetabolizable inducers. A synthesis of 4-thioxylobiose. *Carbohydrate Research* **139**: 123-132.
21. Couchon, N. and A. LeDuy. 1985. A novel process for the production of cellulolytic enzymes. *Biotechnol. Bioeng.* **27**: 456-462.
22. Tangnu, K.S., H.W. Blanch and C.R. Wilke. 1981. Enhanced production of cellulase, hemicellulase, and β -glucosidase by *Trichoderma reesei*. Rut C-30. *Biotechnol. Bioeng.* **23**: 1837-1849.
23. Reese, E.T. and A. Maguire. 1969. Surfactants as stimulants of enzyme production by microorganisms. *Appl. Microbiol.* **17**: 242-245.
24. Szczodrak, J., J. Rogalski and Z. Ilszuk. 1984. Cellulytic activity of moulds. IV. Evaluation of the utility of cellulosic wastes for biosynthesis of cellulases and xylanases by *Aspergillus terreus* F-413. *Acta Microbiol. Polonica.* **3/4**: 217-226.
25. Bisaria, V.S., M. Nanda and K.E. Ghose. 1986. Effect of L(-) sorbose on the release of β -glucosidase by *Trichoderma reesei* QM 9414. *J. Gen. Microbiol.* **132**: 973-978.
26. Biwas, S.R., S.Q. Jana, A.K. Mishra and O. Nanda. 1990. Production, purification and characterization of xylanase from a hyperxylanolytic mutant of *Aspergillus ochraceus*. *Biotechnol. Bioeng.* **35**: 244-251.
27. Warzywoda, M., V. Ferre and J. Pourquie. 1983. Development of culture medium for large-scale production of cellulolytic enzymes by *Trichoderma reesei*. *Biotechnol. Bioeng.* **25**: 3005-3010.
28. Nakajima, T., K. Tsukamoto, T. Watanabe, K. Kainuma and K. Matsuda. 1984. Purification and some properties of an endo-1,4- β -D-xylanase from *Streptomyces* sp. *J. Ferment. Technol.* **62**: 269-276.
29. Ghose, T.K., T. Panda and V.S. Bisaria. 1985. Effect of culture phasing and mannanase on production of cellulase and hemicellulase by mixed culture of *Trichoderma reesei* D 1-6 and *Aspergillus wentii* pt 2804. *Biotechnol. Bioeng.* **27**: 1353-1361.
30. El-Diwany, A.I., A.H.A. Hamdy and M.A. Farid. 1989. Optimization of some culture parameters and properties of β -glucosidase and β -xylosidase from *Penicillium funiculosum* NRRL-13033. *Acta Biotechnol.* **9**: 347-353.

(Received July 27, 1992)